signiert/signed
von: DERMATEST GmbH
am: 08.03.2021
um: 08:48:29 +01

dermatest®
RESEARCH INSTITUTE FOR
RELIABLE RESULTS
TAKING YOUR SUCCESS PERSONALLY

Dermatest® GmbH Engelstr. 37 48143 Münster

Elements of Water GmbH
Unter dem Dostler 4
54293 Trier
Germany

Münster, 05.03.2021

Fachärztliches, dermatologisches Gutachten über einen Epikutantest zur Prüfung auf primäre Irritation und zum Nachweis einer bestehenden Sensibilisierung am Menschen bei einmaliger Applikation des nachstehenden Produktes

Active Anolyte

Auftraggeber: Elements of Water GmbH
Unter dem Dostler 4
54293 Trier
Germany

Testpersonen: 30 Probanden, sämtlich hautgesund

Testkonzentration: unverdünnt

Management System ISO 9001:2015
TÜVRheinland ZERTIFIZIERT
www.tuv.com ID 9105050757

Dermatest® GmbH
Engelstr. 37|48143 Münster
Postfach 2165 |48008 Münster
Germany

info@dermatest.de
www.dermatest.de
Tel. +49 (0) 251 481 637 0
Fax +49 (0) 251 481 637 66

Dr. med. Werner Voss
Geschäftsführer

Amtsgericht Münster HRB 1348

Münsterländische Bank Thie & Co
BLZ 400 300 00 | Konto-Nr. 3 492 784 800
USt-ID: DE126040147 | DE92 3006 0601 0005 3901 50
BIC: MLBK DEH1MUE

PRINZIP UND METHODEN

Das Ziel der Studie ist primäre Hautreizungen durch das Testprodukt und/ oder bestehende Sensibilisierungen gegen das Testprodukt nachzuweisen.

Die Testsubstanzen werden in geeigneten Konzentrationen okklusiv auf die Haut aufgetragen.

Der epikutane Kontakt mit dem Testprodukt ist somit örtlich und zeitlich begrenzt und wird durch die Okklusionsbedingungen intensiviert, sodass die Absorption der Testsubstanzen begünstigt wird. Nach 24, 48 und 72 Stunden wird die Haut untersucht.

Die Okklusion wird verwendet, damit das vermutete topische Allergen leichter durch das Stratum corneum penetrieren kann um die Effektorzellen zu erreichen, die eine lokale Reaktion des Immunsystems hervorrufen könnte.

Damit eine positive Reaktion hervorgerufen werden kann, muss der Schwellenwert der Irritation/ Sensibilität überschritten werden.

Eine positive Reaktion auf einen korrekt applizierten Epikutantest gilt als Nachweis für eine primäre Reizung durch die Testsubstanz, jedoch nicht zwangsläufig als Beweis für eine Sensibilisierung.

Bei bereits bestehenden Sensibilisierungen werden allergische Hautreaktionen durch den Epikutantest provoziert.

DURCHFÜHRUNG

Angehende Testpersonen erhalten eingehende Aufklärung über den Studienverlauf. Wenn eine Teilnahme gewünscht ist und die Studienbedingungen erfüllt sind, unterschreibt der Proband die Einverständniserklärung und die medizinische Anamnese wird erhoben.

20 mg bzw. 20 µl des Testproduktes werden unverdünnt auf ein vorgefertigtes selbstklebendes Pflaster (Curatest® F Folien-Testpflaster, Fa. Lohmann & Rauscher GmbH & Co. KG) auf die klinisch gesunde Haut des oberen Rückens aufgebracht und fixiert. Gewebeartige Produkte werden in einer Fläche von 0,8 cm Ø mit Hilfe des selbstklebenden Pflasters auf den Rücken der Probanden aufgebracht.

Das Testpflaster wird nach einer Expositionsdauer von 24 Stunden abgenommen und das Testareal erstmals dermatologisch und allergologisch beurteilt. Eine zweite und dritte Beurteilung erfolgen nach 48 und 72 Stunden.

Die Bewertung der Testreaktion erfolgt 30 Minuten nach der Entfernung der Testpflaster.

Bei unsicheren Testreaktionen bzgl. ihres Typs (allergisch oder irritativ) werden auch spätere Ablesungen durchgeführt.

Alle Beurteilungen werden bei standardisierter Beleuchtung durchgeführt.

Die Studienteilnehmer werden instruiert die Testareale während der Anwendungsphase trocken zu halten.

PROBANDEN

Die Testgruppe besteht aus 30 erwachsenen, männlichen und weiblichen Probanden.

In dieser Testgruppe sind Probanden mit diverse Hauttypen eingeschlossen, wie zum Beispiel: (sehr) trocken, fettig, misch, normal und sensibel.

EINSCHLUSSKRITERIEN

Personen ab 18 Jahren mit gesunder Haut im Testareal

AUSSCHLUSSKRITERIEN

- Akute organische Krankheit
- Schwangerschaft und Stillzeit
- Bestehende Sensibilisierung auf Inhaltsstoffe des Prüfpflasters
- Schwerwiegende Erkrankungen
- Applikation von wirkstoffhaltigen Präparaten und Pflegemitteln bis 4 Wochen vor Testbeginn
- Einnahme von Medikamenten, die die Hautreaktion beeinträchtigen können Glucocorticoide, Antiallergika, topische Immunmodulatoren, etc.)
- Stark gebräunte Haut

TESTERGEBNISSE

Testkonzentration: Produkt unverdünnt

Nr.	Name	Geschlecht	Alter	Diagnose	24 h	48 h	72 h
1.	AlPe	w	40	hautgesund	-	-	-
2.	BeAn	m	43	hautgesund	-	-	-
3.	BeUr	w	63	hautgesund	-	-	-
4.	DaAn	w	31	hautgesund	-	-	-
5.	DaBu	m	26	hautgesund	-	-	-
6.	DeRo	w	39	hautgesund	-	-	-
7.	FiCl	w	47	hautgesund	-	-	-
8.	FiKa	m	49	hautgesund	-	-	-
9.	FrAn	w	55	hautgesund	-	-	-
10.	FrHe	w	41	hautgesund	-	-	-
11.	FrMa	m	47	hautgesund	-	-	-
12.	GaBo	m	46	hautgesund	?	-	-
13.	GaJo	w	45	hautgesund	-	-	-
14.	GaSt	m	21	hautgesund	-	-	-
15.	GrSa	w	23	hautgesund	-	-	-
16.	HöMa	m	46	hautgesund	-	-	-
17.	JaMo	w	44	hautgesund	-	-	-
18.	JaVi	w	43	hautgesund	-	-	-
19.	KiEl	w	48	hautgesund	-	-	-
20.	KoDi	w	40	hautgesund	-	-	-
21.	KöJo	w	23	hautgesund	-	-	-
22.	MaLi	w	33	hautgesund	-	-	-
23.	PlGi	w	55	hautgesund	-	-	-
24.	PlKe	m	29	hautgesund	-	-	-
25.	PlRo	m	58	hautgesund	-	-	-
26.	RoTa	w	44	hautgesund	-	-	-
27.	ScAl	w	41	hautgesund	-	-	-
28.	SiMi	m	37	hautgesund	-	-	-
29.	SiSi	w	31	hautgesund	-	-	-
30.	SlWi	w	32	hautgesund	-	-	-

TESTERGEBNISSE

WERTUNGEN DES KONTROLLFELDES
Testkonzentration: Leerfeld

Nr.	Name	Geschlecht	Alter	Diagnose	24 h	48 h	72 h
1.	AlPe	w	40	hautgesund	-	-	-
2.	BeAn	m	43	hautgesund	-	-	-
3.	BeUr	w	63	hautgesund	-	-	-
4.	DaAn	w	31	hautgesund	-	-	-
5.	DaBu	m	26	hautgesund	-	-	-
6.	DeRo	w	39	hautgesund	-	-	-
7.	FiCl	w	47	hautgesund	-	-	-
8.	FiKa	m	49	hautgesund	-	-	-
9.	FrAn	w	55	hautgesund	-	-	-
10.	FrHe	w	41	hautgesund	-	-	-
11.	FrMa	m	47	hautgesund	-	-	-
12.	GaBo	m	46	hautgesund	-	-	-
13.	GaJo	w	45	hautgesund	-	-	-
14.	GaSt	m	21	hautgesund	-	-	-
15.	GrSa	w	23	hautgesund	-	-	-
16.	HöMa	m	46	hautgesund	-	-	-
17.	JaMo	w	44	hautgesund	-	-	-
18.	JaVi	w	43	hautgesund	-	-	-
19.	KiEl	w	48	hautgesund	-	-	-
20.	KoDi	w	40	hautgesund	-	-	-
21.	KöJo	w	23	hautgesund	-	-	-
22.	MaLi	w	33	hautgesund	-	-	-
23.	PlGi	w	55	hautgesund	-	-	-
24.	PlKe	m	29	hautgesund	-	-	-
25.	PlRo	m	58	hautgesund	-	-	-
26.	RoTa	w	44	hautgesund	-	-	-
27.	ScAl	w	41	hautgesund	-	-	-
28.	SiMi	m	37	hautgesund	-	-	-
29.	SiSi	w	31	hautgesund	-	-	-
30.	SlWi	w	32	hautgesund	-	-	-

AUSWERTUNGSKRITERIEN

Die Beurteilung erfolgt aufgrund der morphologischen Ausprägung modifiziert nach Empfehlungen der ICDRG (Fregert S (1981/ 2nd edition) Manual of Contact Dermatitis. On behalf of the International Contact Dermatitis Research Group and the North American Contact Dermatitis Group, Munksgaard Publishers, Copenhagen).

Tabelle 3: Beurteilung von Epikutantestreaktionen

Symbol	**Morphe**	**Bedeutung**
-	keine Reaktion	negativ
?	nur Erythem, kein Infiltrat	fraglich
+	Erythem, Infiltrat, evtl. diskrete Papeln	einfach-positive Reaktion
++	Erythem, Infiltrat, Papeln, Vesikel	zweifach-positive Reaktion
+++	Erythem, Infiltrat, konfluierende Vesikel	dreifach-positive Reaktion
ir	versch. Veränd. (Seifeneffekt, Vesikel, Blase, Nekrose)	irritativ
nt		nicht getestet

ALLGEMEINE DERMATOLOGISCHE INTERPRETATIONSKRITERIEN:

Wichtig ist es bei der Beurteilung der Ergebnisse, zwischen toxischen und allergischen Reaktionen zu unterscheiden. In der Regel werden die zum Zeitpunkt 72 Std. oder später als "+" bis "+++" beurteilten Reaktionen als "allergisch" gedeutet.

In die Bewertung kann die Reaktionsdynamik einfließen. Testreaktionen allergischen Typs können bestehen oder sogar stärker werden am Tag nach Entfernung der Testpflaster ("Plateau-Muster" bzw. "Crescendo"). Im Gegensatz dazu spricht ein "Decrescendo"-Muster (abschwächen der Reaktion nach Entfernen des Pflasters) eher für eine irritative Reaktion.

Bei Spätreaktionen, die etwa 10-14 Tage nach Applikation des Epikutantests erstmalig auftreten, ist an eine ("iatrogene") Sensibilisierung durch den Test zu denken.

Eine irritative/ allergische Reaktion äußert sich in einem Erythem und kann auch ein leichtes Infiltrat aufweisen.

Papeln, Vesikeln und Blasen können sowohl als Zeichen einer Irritation als auch einer Allergie auftreten, wohingegen Pusteln und Nekrosen starke irritative Reaktionen kennzeichnen.

Das ursprüngliche Applikationsareal kann in beiden Reaktionsformen deutlich überschritten werden. Zudem kann die individuelle Ausprägung der Reaktion stark schwanken.

BEURTEILUNG DER TESTERGEBNISSE

Bei 29 der 30 Probanden kam es bei den Epikutantestungen nach internationalen Richtlinien der ICDRG (International Contact Dermatitis Research Group) nach 24, 48 und 72 Stunden zu keinen Hautveränderungen im Testbezirk.

Nach 24 Stunden zeigte sich bei einem Probanden eine milde Reaktion, die nach 48 Stunden nicht mehr sichtbar war.

Vereinzelte, milde Reaktionen im Epikutantest sind in einer Testgruppe normal und im Rahmen der Prüfbedingungen im Gegensatz zur bestimmungsgemäßen Anwendung zu sehen. Dennoch könnte die Anwendung des Produktes bei empfindlicher Haut zu milden, vorübergehenden Hautreaktionen führen.

Dermatest GmbH
Engelstrasse 37
48143 Münster Germany
dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
Taking your success personally
Fon: +49-(0)251-4816370
Fax: +49-(0)251-48163766
www.dermatest.de

Dr. med. Gerrit Schlippe
Facharzt für Dermatologie,
Venerologie

Dr. med. Werner Voss
Facharzt für Dermatologie,
Venerologie, Allergologie,
Phlebologie und Umweltmedizin

Literatur:

1. Suzuki, J., Environ Dermatol 4-3:202-21-1997
2. ICDRG, Proposal for a revised international standard series of patch tests, Contact Dermatitis, No. 36, 121-123 (1997)
3. U.S. Department of Health and Human Services Food and Drug Administration, April 1999 http:/www.fda.gov/cber/guidelines.htm
4. Scientific Basis of Patch Testing – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 2, 43-50 (2002)
5. Scientific Basis of Patch Testing Part II – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 3, 91-96 (2002)
6. Scientific Basis of Patch Testing Part III – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 4, 131-133 (2002)
7. Cosmetics & Toiletries magazine, www.CosmeticsandToiletries.com, Vol. 127, No. 5/May 2012, pages 356-360

dermatest®
RESEARCH INSTITUTE FOR
RELIABLE RESULTS
TAKING YOUR SUCCESS PERSONALLY

Dermatest® GmbH Engelstr. 37 48143 Münster

Elements of Water GmbH
Unter dem Dostler 4
54293 Trier
Germany

Muenster, 05.03.2021

Dermatological report on human Patch Test for primary skin irritation and to detect existing sensitisations of human subjects after single application of

Active Anolyte

Customer:	Elements of Water GmbH Unter dem Dostler 4 54293 Trier Germany
Test Panel:	30 panellists of either sex, all without visible skin diseases or known hypersensitivity
Concentration of the product:	undiluted

Management System ISO 9001:2015
TÜVRheinland ZERTIFIZIERT
www.tuv.com ID 9105050757

PRINCIPLE AND METHODS

The objective of the study is to detect primary skin irritation potential and/ or existing allergic sensitisation to the test substance.

The test substance is applied to the skin of the panellist via an occlusive patch at a suitable concentration.

The patch limits contact of the panellist's skin with the test substance to a local area and exposure is exaggerated due to the occlusive conditions. The skin is checked at 24, 48 and 72 hours.

The occlusion eases the absorption of the suspected topical allergen allowing it to penetrate the stratum corneum to the viable (effector) cells of the skin and thus presenting a local challenge to the immune system.

If the threshold level of sensitivity is reached, a positive reaction could potentially be induced.

A positive reaction to a correctly applied patch provides evidence of primary irritation to the substance tested, but is not necessarily evidence of sensitisation.

Patch testing provokes allergic skin reactions in already sensitised panellists.

PROCEDURES

Prospective panellists receive a complete explanation of study procedures. If they wish to participate and agree to the conditions of the study, panellists sign a written, informed consent and provide a medical history.

20 mg/ 20 µl of the undiluted test product is applied to an adhesive plaster (Curatest® F Folien-Testpflaster, Fa. Lohmann & Rauscher GmbH & Co. KG) and affixed to clinically healthy skin on the upper back. Textile products are affixed with a sample size of 0,8 cm Ø with the adhesive plaster on the upper back.

After a 24 hour exposure period, the plaster is removed and the exposed skin is dermatologically assessed and graded. The second and third assessments are performed after 48 and 72 hours respectively.

All assessments are conducted 30 minutes after removal of the test plaster.

Where a positive reaction is observed, but it is unclear whether the observed reaction is due to sensitisation or irritation, subsequent readings can be performed.

All assessments are performed under standard lighting conditions.

The panellists are instructed to keep the test sites dry.

PANELLISTS

The test panel included 30 adult male and female subjects.

This test group includes test persons with various skin types, such as: (very) dry, oily, mixed, normal and sensitive.

INCLUSION CRITERIA

Subjects aged 18 years and above with healthy skin in the test area

EXCLUSION CRITERIA

- Acute diseases
- Pregnancy and lactation period
- Sensitisation to ingredients of the test plaster
- Severe illnesses
- Application of pharmaceutical products and skin care products with active ingredients until 4 weeks before testing
- Intake of drugs that possibly can interfere with skin reactions (steroids, antiallergics, topical immuno modulator, etc.)
- Extremely tanned skin

RESULTS

Table 1: Results of patch testing for the test substance
Concentration of the product: undiluted

No.	Name	Gender	Age	Diagnosis	24 h	48 h	72 h
1.	AlPe	f	40	healthy skin	-	-	-
2.	BeAn	m	43	healthy skin	-	-	-
3.	BeUr	f	63	healthy skin	-	-	-
4.	DaAn	f	31	healthy skin	-	-	-
5.	DaBu	m	26	healthy skin	-	-	-
6.	DeRo	f	39	healthy skin	-	-	-
7.	FiCl	f	47	healthy skin	-	-	-
8.	FiKa	m	49	healthy skin	-	-	-
9.	FrAn	f	55	healthy skin	-	-	-
10.	FrHe	f	41	healthy skin	-	-	-
11.	FrMa	m	47	healthy skin	-	-	-
12.	GaBo	m	46	healthy skin	-	-	-
13.	GaJo	f	45	healthy skin	**?**	-	-
14.	GaSt	m	21	healthy skin	-	-	-
15.	GrSa	f	23	healthy skin	-	-	-
16.	HöMa	m	46	healthy skin	-	-	-
17.	JaMo	f	44	healthy skin	-	-	-
18.	JaVi	f	43	healthy skin	-	-	-
19.	KiEl	f	48	healthy skin	-	-	-
20.	KoDi	f	40	healthy skin	-	-	-
21.	KöJo	f	23	healthy skin	-	-	-
22.	MaLi	f	33	healthy skin	-	-	-
23.	PlGi	f	55	healthy skin	-	-	-
24.	PlKe	m	29	healthy skin	-	-	-
25.	PlRo	m	58	healthy skin	-	-	-
26.	RoTa	f	44	healthy skin	-	-	-
27.	ScAl	f	41	healthy skin	-	-	-
28.	SiMi	m	37	healthy skin	-	-	-
29.	SiSi	f	31	healthy skin	-	-	-
30.	SlWi	f	32	healthy skin	-	-	-

RESULTS

Table 2: RESULTS of patch testing for the CONTROL
Concentration of the product: blank patch test

No.	Name	Gender	Age	Diagnosis	24 h	48 h	72 h
1.	AlPe	f	40	healthy skin	-	-	-
2.	BeAn	m	43	healthy skin	-	-	-
3.	BeUr	f	63	healthy skin	-	-	-
4.	DaAn	f	31	healthy skin	-	-	-
5.	DaBu	m	26	healthy skin	-	-	-
6.	DeRo	f	39	healthy skin	-	-	-
7.	FiCl	f	47	healthy skin	-	-	-
8.	FiKa	m	49	healthy skin	-	-	-
9.	FrAn	f	55	healthy skin	-	-	-
10.	FrHe	f	41	healthy skin	-	-	-
11.	FrMa	m	47	healthy skin	-	-	-
12.	GaBo	m	46	healthy skin	-	-	-
13.	GaJo	f	45	healthy skin	-	-	-
14.	GaSt	m	21	healthy skin	-	-	-
15.	GrSa	f	23	healthy skin	-	-	-
16.	HöMa	m	46	healthy skin	-	-	-
17.	JaMo	f	44	healthy skin	-	-	-
18.	JaVi	f	43	healthy skin	-	-	-
19.	KiEl	f	48	healthy skin	-	-	-
20.	KoDi	f	40	healthy skin	-	-	-
21.	KöJo	f	23	healthy skin	-	-	-
22.	MaLi	f	33	healthy skin	-	-	-
23.	PlGi	f	55	healthy skin	-	-	-
24.	PlKe	m	29	healthy skin	-	-	-
25.	PlRo	m	58	healthy skin	-	-	-
26.	RoTa	f	44	healthy skin	-	-	-
27.	ScAl	f	41	healthy skin	-	-	-
28.	SiMi	m	37	healthy skin	-	-	-
29.	SiSi	f	31	healthy skin	-	-	-
30.	SlWi	f	32	healthy skin	-	-	-

INTERPRETATION CRITERIA

The assessment is based on the morphologic changes detailed in the modified guidelines of ICDRG (Fregert S (1981/ 2nd edition) Manual of Contact Dermatitis. On behalf of the International Contact Dermatitis Research Group and the North American Contact Dermatitis Group, Munksgaard Publishers,Copenhagen).

Table 3: Grading of the patch test reactions

Symbol	**Morphology**	**Meaning**
-	no reaction	negative
?	only erythema, no infiltration	doubtful
+	erythema, infiltration, possibly discrete papules	weak positive reaction
++	erythema, infiltration, papules, vesicles	strong positive reaction
+++	erythema, infiltration, papules, confluent vesicles	extreme severe positive reaction
ir	different changes (soap effect, vesicles, bulla, necrosis)	irritative
nt		not tested

GENERAL DERMATOLOGICAL INTERPRETATION CRITERIA:

The distinction between irritation and allergy is of importance. As a general rule, a positive reaction is said to be „allergic“ if it has been graded as “+” to “+++ “ up to 72 hours or beyond.

Understanding the dynamics of the reaction may aid the assessment.

Allergic test reactions could persist ("Plateau-type”) or even worsen ("Crescendo-type") on the day after the plaster has been removed). A “Decrescendo”-type (decrease of reaction after removal of plaster) on the other hand, indicates irritation.

If delayed reactions only develop 10-14 days after application, (“iatrogenic”) sensitisation should be considered.

Irritative and allergic reactions present erythema and could also cause infiltration.

Papules, vesicles and bullae could demonstrate irritation as well as allergy, whereas pustules and necrosis point to severe irritation reactions.

Both reactions could spread beyond the original application site.

Moreover the individual expression of a reaction lies within a wide range.

CONCLUSION

For 29 of 30 panellists no evidence of any skin disorder was detected in the test area of any of the 30 panellists after conducting patch testing for 24, 48 and 72 hours according to the internationally recognised guidelines of ICDRG (International Contact Dermatitis Research Group).

After 24 hours, one subject showed a mild reaction that was no longer visible after 48 hours.

Sporadic, mild reactions in the epicutaneous test are normal in a test group and must be seen in the context of the test conditions in contrast to the intended use. Still, use of the product could cause mild, transient skin reactions in sensitive skin.

Dermatest GmbH
Engelstrasse 37
48143 Münster Germany
dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
Taking your success personally
Fon: +49-(0)251-4816370
Fax: +49-(0)251-48163766
www.dermatest.de

Dr. med. Gerrit Schlippe
Investigating specialist
for dermatology, venereology

Dr. med. Werner Voss
Investigating specialist
for dermatology, allergology,
venereology, phlebology
and environmental medicine

Literature:

1. Suzuki, J., Environ Dermatol 4-3:202-21-1997
2. ICDRG, Proposal for a revised international standard series of patch tests, Contact Dermatitis, No. 36, 121-123 (1997)
3. U.S. Department of Health and Human Services Food and Drug Administration, April 1999 http:/www.fda.gov/cber/guidelines.htm
4. Scientific Basis of Patch Testing – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 2, 43-50 (2002)
5. Scientific Basis of Patch Testing Part II – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 3, 91-96 (2002)
6. Scientific Basis of Patch Testing Part III – S. Iris Ale and Howard I. Maibach, Dermatol. Beruf Umwelt / Occup. Environ. Dermatol. 50, Nr. 4, 131-133 (2002)
7. Cosmetics & Toiletries magazine, www.CosmeticsandToiletries.com, Vol. 127, No. 5/May 2012, pages 356-360

dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
TAKING YOUR SUCCESS PERSONALLY

Dermatest® GmbH Engelstr. 37 48143 Münster

Elements of Water GmbH
Unter dem Dostler 4
54293 Trier
Germany

Münster, 05.03.2021

Zertifikat

über das Produkt

Active Anolyte

Dermatologische Prüfungen am Menschen 2021

Studiennummer: 2102190112

Die von mir unter fachärztlicher Kontrolle durchgeführten dermatologischen Prüfungen Ihres o. g. Produktes bestand das Produkt mit

„sehr gut“

Bei dem getesteten Präparat traten im Epikutantest nach internationalen Richtlinien bei weniger als 10% der Probanden sehr milde vorübergehende Reaktionen auf. Bei allen anderen Probanden traten k e i n e toxisch- irritativen Unverträglichkeitsreaktionen auf.

Vereinzelte, milde Reaktionen im Epikutantest sind in einer Testgruppe normal und im Rahmen der Prüfbedingungen im Gegensatz zur bestimmungsgemäßen Anwendung zu sehen. Somit ist das Produkt nicht beschränkt in der Bewertung „dermatologisch getestet“.

Dr. med. Gerrit Schlippe
Facharzt für Dermatologie, Venerologie

Dermatest GmbH
Engelstrasse 37
48143 Münster Germany
dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
Taking your success personally
Fon: +49-(0)251-4816370
Fax: +49-(0)251-48163766
www.dermatest.de

Dr. med. Werner Voss
Facharzt für Dermatologie, Venerologie, Allergologie, Phlebologie und Umweltmedizin

TÜVRheinland ZERTIFIZIERT
Management System ISO 9001:2015
www.tuv.com ID 9105060763

dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
TAKING YOUR SUCCESS PERSONALLY

Dermatest® GmbH Engelstr. 37 48143 Münster

Elements of Water GmbH
Unter dem Dostler 4
54293 Trier
Germany

Muenster, 05.03.2021

Certificate

for the Product

Active Anolyte

Dermatological test on humans in 2021

Study number: 2102190112

The dermatological test performed by us on your product under the control of dermatological specialists was passed for this product with the rating of

„excellent“

Less than 10% of the test persons showed very mild transient reactions in the epicutaneous test according to international guidelines with the tested preparation. For all other subjects this product did not lead to any toxic-irritative intolerance reactions.

Sporadic, mild reactions in the epicutaneous test are normal in a test group and must be seen in the context of the test conditions in contrast to the intended use. Thus, the product is not restricted in the evaluation "dermatologically tested".

Dr. med. Gerrit Schlippe
Investigating specialist
for dermatology, venereology

Dermatest GmbH
Engelstrasse 37
48143 Münster Germany
dermatest®
RESEARCH INSTITUTE FOR RELIABLE RESULTS
Taking your success personally
Fon: +49-(0)251-4816370
Fax: +49-(0)251-48163766
www.dermatest.de

Dr. med. Werner Voss
Investigating specialist
for dermatology, allergology,
venerology, phlebology
and environmental medicine

TÜVRheinland ZERTIFIZIERT
Management System ISO 9001:2015
www.tuv.com ID 9105060753